

AD610593

EXPONENTIAL LIFE TEST PROCEDURES  
WHEN THE DISTRIBUTION HAS  
MONOTONE FAILURE RATE

by

Richard E. Barlow

Frank Proschan

ORC 64-29  
OCTOBER 1964




OPERATIONS RESEARCH CENTER

INSTITUTE OF ENGINEERING RESEARCH

ADDITIONAL COPY

UNIVERSITY OF CALIFORNIA - BERKELEY



ERRATA SHEET for "Exponential Life Test Procedures When the Distribution has Monotone Failure Rate," by R. E. Barlow and Frank Proschan.

Page 11, line 10 and page 12, line 7: Replace by:

$$R_1(t) = \begin{cases} (1 - \frac{t}{Z_r})^{r-1} & \text{if } t < Z_r \\ 0 & \text{otherwise} \end{cases}$$

Page 12, line 8: Replace by:

$$\geq \begin{cases} 1 - \frac{t}{\phi(Z_r)} & \text{if } t < Z_r \\ 0 & \text{otherwise.} \end{cases}$$

Page 13, line 6: Replace "Theorem 2.1" by "Corollary 2.2".

Page 13, line 12: Insert "(larger)" after "smaller".

Page 13, 2nd line from bottom; and page 32, line 5: Replace " $\geq$ " by " $\geq(\leq)$ ".

Page 13, bottom line: Replace " $\leq E$ " by " $\leq(\geq) E$ ".

Page 16, line 3: Insert "<" between "T" and "t".

Page 22, line 3: Put subscript " $\Delta$ " on "G" and " $\bar{G}$ ".

Page 22, bottom line:

$$\text{Replace: } e^{x \log \frac{q}{\xi_p}} \text{ by } e^{\frac{x \log q}{\xi_p}}.$$

Page 30, line 14: Replace by " $\bar{H}_1(x)$ ".

Page 31, line 5: Insert "(DFR)" after "IFR".

EXPONENTIAL LIFE TEST PROCEDURES  
WHEN THE DISTRIBUTION HAS MONOTONE FAILURE RATE

By

Richard E. Barlow, University of California, Berkeley,  
Frank Proschan, University of California, Berkeley, and  
Boeing Scientific Research Laboratories.

October 1964

ORC 64-29

This research has been partially supported by the Office of Naval Research under Contract Nomr-3656(18) with the University of California. The report will also be issued by Boeing Scientific Laboratories. Reproduction in whole or in part is permitted for any purpose of the United States Government.

## SUMMARY

A number of estimates and tests for mean life and other parameters derived under the exponential distribution assumption are studied under the alternative condition that the distribution has increasing (decreasing) failure rate. The estimates considered are, for the most part, based on censored and truncated samples. It is shown that these estimates generally favor the producer (consumer) in the IFR (DFR) case. Properties of order statistics and their spacings from distributions with increasing (decreasing) failure rate are presented.

1. Introduction. In a fundamental paper in the literature of life testing Epstein and Sobel (1953) introduce life test procedures based on the exponential distribution. These procedures have been codified in a Department of Defense handbook (1960) and are now widely employed. Zelen and Dannemiller (1961) show by sampling from Weibull distribution alternatives that these procedures are not robust in testing for mean life. However, as Antelman and Savage (unpub.) have pointed out they may be robust in testing for certain percentiles. For certain loss functions based on percentiles, these procedures seem to be robust. Since statistical procedures based on the exponential distribution have a great deal of intuitive appeal and computational simplicity we investigate their properties relative to alternative distributions having increasing failure rate (IFR) or decreasing failure rate (DFR).

This paper essentially confirms, theoretically and more generally, the sampling results of the Zelen-Dannemiller paper for statistics derived under the exponential assumption. Using Weibull distribution alternatives (with parameter values which insure that the distribution has increasing failure rate) Zelen-Dannemiller show that the use of these statistics may result in substantially increasing the probability of accepting items having poor mean lives. We show that these estimates for the mean are positively (negatively) biased when the distribution is IFR (DFR). Also we obtain bounds on the expected values of the exponential estimates for the distribution function and bounds on the expected values of the order statistics.

In the last section various properties of IFR (DFR) order statistics are presented.

Preliminaries. Let  $X$  denote a random variable with right continuous distribution  $F$  such that  $F(0^-) = 0$ . If  $F$  has density  $f$  then  $r(t) = \frac{f(t)}{[1 - F(t)]}$  is known as the failure rate. Note that  $r(t) = -\frac{d}{dt} \log[1 - F(t)]$  when a density exists. For this reason, we say that  $F$  is IFR (DFR) for increasing (decreasing) failure rate if  $\log[1 - F(t)]$  is concave where finite (convex on  $[0, \infty)$ ). Note that any IFR (DFR) distribution with specified mean can be expressed as the limit of continuous IFR (DFR) distributions with the same mean. Hence for many of our results it is sufficient to confine attention to continuous IFR (DFR) distributions.

We often use the well known fact that if  $F$  is continuous, then  $Y = -\theta \log \bar{F}(X)$  is exponentially distributed with mean  $\theta$  where  $\bar{F}(x) = 1 - F(x)$ . Repeatedly we use the fact that if  $F$  is IFR with mean  $\theta$  then there exists  $x_0 \geq \theta$  such that

$$y = -\theta \log \bar{F}(x) \begin{cases} < x & \text{for } x < x_0 \\ \geq x & \text{for } x \geq x_0 \end{cases}.$$

This is evident from log concavity and the bounds on IFR distributions given in Barlow and Marshall (1964). The inequalities are reversed when  $F$  is DFR.

Unless otherwise indicated we denote ordered observations from a random sample of size  $n$  based on a random variable  $X$

by  $X_1 \leq \dots \leq X_n$ . We define  $X_0 = 0$ .

2. Estimates based on censored samples. Assume  $n$  items are put on life test and let  $X_1 \leq X_2 \leq \dots \leq X_n$  denote the ordered observations. If  $F$  has density  $f$  such that

$$f(x; \theta) = \begin{cases} \frac{1}{\theta} e^{-\frac{x}{\theta}} & x \geq 0 \\ 0 & x < 0 \end{cases}$$

then

$$(2.1) \quad \hat{\theta}_{r,n} = \frac{\sum_{i=1}^r X_i + (n-r)X_r}{r} = \frac{\sum_{i=1}^r (n-i+1)(X_i - X_{i-1})}{r}$$

$(1 \leq r \leq n)$  is the maximum likelihood and minimum variance unbiased estimate for  $\theta$  based on the first  $r$  order statistics (Epstein and Sobel, 1953).

The normalized spacings  $\bar{D}_i = (n-i+1)(X_i - X_{i-1})$  which enter into the computation of  $\hat{\theta}_{r,n}$  have a natural intuitive appeal. They have also been used as the basis for a statistic to test for IFR (Proschan and Pyke, in preparation). We shall derive and use several properties of these spacings when  $F$  is IFR (DFR). Since the normalized spacings are independent and identically distributed in the exponential case (Epstein and Sobel, 1953), Theorem 2.1 below is quite intuitive.

A random variable  $X$  is said to be stochastically smaller than a random variable  $Y$  if and only if  $P[X \geq x] \leq P[Y \geq x]$  for all  $x$ .

Theorem 2.1. If  $F$  is IFR (DFR) the normalized spacings  $(n - i + 1)(X_i - X_{i-1})$  are stochastically decreasing (increasing) in  $i$ .

Proof. Assume  $F$  is IFR and let  $\bar{F}(x) = 1 - F(x)$ . Note that

$$P[nX_1 \geq x] = [\bar{F}(\frac{x}{n})]^n \geq [\bar{F}(\frac{x}{n-1})]^{n-1}$$

since  $[\bar{F}(t)]^{\frac{1}{t}}$  is decreasing in  $t$ . Let

$$F_u(x) = \frac{F(u+x) - F(u)}{\bar{F}(u)}$$

and note  $F_u(x) \geq F(x)$ . Given that  $X_1 = u$  is observed,  $X_2 - X_1$  is distributed as the first order statistic from a sample of size  $n - 1$  each with distribution  $F_u(x)$ . Hence

$$P[(n-1)(X_2 - X_1) \geq x \mid X_1 = u] = [\bar{F}_u(\frac{x}{n-1})]^{n-1}.$$

Conditioning on  $X_1$  we have

$$\begin{aligned} P[nX_1 \geq x] &= [\bar{F}(\frac{x}{n})]^n \geq [\bar{F}_u(\frac{x}{n-1})]^{n-1} \\ &= P[(n-1)(X_2 - X_1) \geq x \mid X_1 = u] \end{aligned}$$

for all  $u \geq 0$ . Unconditioning

$$\begin{aligned} P[nX_1 \geq x] &\geq \int_0^\infty [\bar{F}_u(\frac{x}{n-1})]^{n-1} dG(u) \\ &= P[(n-1)(X_2 - X_1) \geq x] \end{aligned}$$

where  $G(u) = 1 - [\bar{F}(u)]^n$  is the distribution of  $X_1$ . Hence we have shown that  $nX_1$  is stochastically larger than  $(n-1)(X_2 - X_1)$ .

In a similar manner we can show that  $(n - i + 1)(X_i - X_{i-1})$  is stochastically larger than  $(n - i)(X_{i+1} - X_i)$  for  $i = 2, 3, \dots, n$ .

All inequalities are reversed for DFR distributions. ||

As an immediate consequence of Theorem 2.1 we have that



$$E \left\{ [(n-i+1)(X_i - X_{i-1})]^\alpha \right\}$$

is decreasing (increasing) in  $i$  for  $\alpha \geq 0$  when  $F$  is IFR (DFR). Using this fact we can show that  $\hat{\theta}_{r,n}$  is positively biased when  $F$  is IFR.

Corollary 2.2. If  $F$  is IFR with mean  $\theta$ , then

$$\theta \leq E[\hat{\theta}_{r,n}] \leq n \frac{\theta}{r} \quad \text{for } r = 1, 2, \dots, n.$$

All inequalities are sharp.

Proof. From Barlow and Proschan (1964a, p.33) we know that

$$E[\hat{\theta}_{1,n}] \geq \theta. \quad \text{Also} \quad h(r) = \sum_{i=1}^r \left\{ E[(n-i+1)(X_i - X_{i-1})] - \theta \right\}$$

exhibits at most one sign change as a function of  $r$  since

$$E[(n-i+1)(X_i - X_{i-1})]$$

is decreasing in  $i$  by Theorem 2.1. But  $h(1) \geq 0$  and  $h(n) = 0$ ,

which implies  $h(r) \geq 0$  for  $r = 1, 2, \dots, n$ . Hence

$$E[\hat{\theta}_{r,n}] \geq \theta.$$

Clearly the bound is attained by the exponential distribution so that it is sharp.

To show the upper bound we note

$$\sum_{i=1}^r X_i + (n-r) X_r \leq \sum_{i=1}^n X_i$$

for every sample realization. Hence

$$E[r \hat{\theta}_{r,n}] \leq n \theta$$

or 
$$E[\hat{\theta}_{r,n}] \leq n \frac{\theta}{r}.$$

Since equality is attained with distributions degenerate at  $\theta$  (which is the limit of IFR distributions) the bound is sharp. ||

Corollary 2.3. If  $F$  is DFR with mean  $\theta$ , then

$$0 \leq E[\hat{\theta}_{r,n}] \leq \theta \quad \text{for } 1 \leq r \leq n.$$

All inequalities are sharp.

Proof. The upper bound follows from Theorem 2.1 and the method of proof in Corollary 2.2. To show that the lower bound is sharp, let

$$F(x) = \begin{cases} 0 & x < 0 \\ e^{-\frac{\epsilon x}{\theta}} & x \geq 0 \end{cases}$$

where  $\epsilon > 0$  is arbitrary. Then  $F$  is DFR with mean  $\theta$  and

$$\begin{aligned} P[X_i \geq x] &= \sum_{j=0}^{i-1} \binom{n}{j} [F(x)]^j [1-F(x)]^{n-j} \\ &\leq \sum_{j=0}^{i-1} \binom{n}{j} e^{-\frac{\epsilon x j}{\theta}} = e^{-\frac{\epsilon x (i-1)}{\theta}} \end{aligned}$$

Hence

$$E[X_i] = \int_0^{\infty} P[X_i \geq x] dx \leq \sum_{j=0}^{i-1} \binom{n}{j} \frac{\epsilon^{n-j-1} \theta}{(n-j)} < 2^n \epsilon \theta$$

when  $0 \leq \epsilon < 1$ . Since  $\epsilon$  is arbitrary we see that

$$E[X_i] \geq 0 \quad (1 \leq i \leq n)$$

is sharp. ||

For convenience we now denote the  $i^{\text{th}}$  order statistic from a sample of size  $n$  by  $X_{i,n}$ .

Theorem 2.4. If  $F$  is IFR (DFR)  $(n-i+1)(X_{i,n} - X_{i-1,n})$  is stochastically increasing (decreasing) in  $n$  for fixed  $i$ . Hence

$$E[\hat{\theta}_{r,n}] \underset{(\geq)}{\leq} E[\hat{\theta}_{r,n+1}] \quad (1 \leq r \leq n)$$

Proof. Assume  $F$  is IFR. Let  $G_{i,n}(x) = P[X_{i,n} \leq x]$  and note that  $G_{i,n}(x) \leq G_{i,n+1}(x)$  for samples from any distribution.

Now

$$\begin{aligned} P[(n-i)(X_{i+1,n} - X_{i,n}) \geq x] &= \int_0^\infty \left[ F_u\left(\frac{x}{n-i}\right) \right]^{n-i} dG_{i,n}(u) \\ &\leq \int_0^\infty \left[ F_u\left(\frac{x}{n+1-i}\right) \right]^{n+1-i} dG_{i,n}(u) \\ &\leq \int_0^\infty \left[ F_u\left(\frac{x}{n+1-i}\right) \right]^{n+1-i} dG_{i,n+1}(u) \\ &= P[(n+1-i)(X_{i+1,n+1} - X_{i,n+1}) \geq x]; \end{aligned}$$

the first inequality holds since  $[F(t)]^{\frac{1}{t}}$  is decreasing in  $t$  when  $F$  is IFR.

All inequalities are reversed when  $F$  is DFR. ||

Thus when  $F$  is IFR, the estimate,  $\hat{\theta}_{r,n}$ , of mean life based on a sample censored on the right becomes worse with increasing  $n$  when  $r$  ( $1 \leq r \leq n$ ) remains fixed.

Acceptance Sampling. Statistical methods for testing hypotheses about the mean of an exponential distribution depend on the statistic,  $\hat{\theta}_{r,n}$ , in the case of censored samples (Epstein, 1960a). For testing the hypothesis  $H_0: \theta = \theta_0$  against the alternative  $H_1: \theta = \theta_1 < \theta_0$  subject to  $P[\text{reject } \theta = \theta_0 \mid \theta_0 \text{ true}] = \alpha$ , the rejection region is of the form

$$\hat{\theta}_{r,n} \leq \frac{\theta_0 \chi_{1-\alpha}^2(2r)}{2r}.$$

If  $\chi_{1-\alpha}^2(2r) < 2r$ , then we shall prove, using Lemma 2.5, that

$$\begin{aligned}
 (2.2) \quad & P \left[ \hat{\theta}_{r,n} \leq \frac{\theta_0 x_{1-\alpha}^2(2r)}{2r} \mid F \text{ IFR}, \theta = \theta_0 \right] \\
 & \leq P \left[ \hat{\theta}_{r,n} \leq \frac{\theta_0 x_{1-\alpha}^2(2r)}{2r} \mid F(x) = 1 - e^{-\frac{x}{\theta_0}} \right] = \alpha
 \end{aligned}$$

so that the exponential test provides a size  $\alpha$  test when the failure distribution is IFR. To see (2.2) we need the following easily verified result which we present without proof.

Lemma 2.5. If  $\phi$  is concave  $\phi(0) = 0$ , and  $a_i \geq 1$ ,  $x_i \geq 0$  ( $i = 1, 2, \dots, n$ ), then

$$\phi \left[ \sum_{i=1}^n a_i x_i \right] \leq \sum_{i=1}^n a_i \phi(x_i)$$

Let  $\phi^{-1}(y) = -\theta \log \bar{F}(y)$  so that  $X_i = \phi(Y_i)$

where  $Y_i$  is the  $i^{\text{th}}$  order statistic of an exponentially distributed random variable with mean  $\theta$ . Then

$$\begin{aligned}
 r\hat{\theta}_{r,n} &= \sum_{i=1}^r \phi(Y_i) + (n-r)\phi(Y_r) \\
 &\geq \phi \left[ \sum_{i=1}^r Y_i + (n-r)Y_r \right]
 \end{aligned}$$

by Lemma 2.5. Using the bounds on IFR distributions (Barlow

and Marshall, 1964) and letting  $Z_r^* = \sum_{i=1}^r Y_i + (n-r)Y_r$ , we have

$$P_\theta [\phi(Z_r^*) \leq c] \leq P_\theta [Z_r^* \leq c]$$

when  $c < \theta$ . We obtain (2.2) by letting  $\theta = \theta_0$  and

$$c = \frac{\theta_0 \chi_{1-\alpha}^2(2r)}{2}.$$

Sampling with Replacement. Suppose now that failed items are replaced at failure. In this case the bias of the usual estimate for  $\theta$  is even greater than in the non-replacement case.

Let  $X_i^*$  denote the time of the  $i^{\text{th}}$  failure when failed items are replaced. The maximum likelihood estimate for  $\theta$  based on the exponential assumption is, in this case

$$\begin{aligned}\hat{\theta}_{r,n}^* &= \frac{1}{r} [nX_1^* + n(X_2^* - X_1^*) + \dots + n(X_r^* - X_{r-1}^*)] \\ &= \frac{nX_r^*}{r}\end{aligned}$$

(Epstein, 1960b).

Theorem 2.5. If  $F$  is IFR with mean  $\theta$ , then

$$\theta \leq E[\hat{\theta}_{r,n}^*] \leq E[\hat{\theta}_{r,n}^{**}] \leq \frac{n}{r} E[X_r] \quad 1 \leq r \leq n.$$

Proof. Clearly  $X_r^* \leq X_r$  for any distribution  $F$  so that the upper bound is obvious. To show the remaining inequality we introduce the following fictitious replacement policy:

Policy A: Replace a failed item with a good item of the same "age".

Let  $X_i^{**}$  denote the time of the  $i^{\text{th}}$  failure under this policy and

$$\hat{\theta}_{r,n}^{**} = \frac{1}{r} \sum_{i=1}^r (n - i + 1)(X_i^{**} - X_{i-1}^{**}).$$

It is clear that  $E[\hat{\theta}_{r,n}^{**}] \leq E[\hat{\theta}_{r,n}^*]$

since under the IFR assumption the conditional mean life of an aged item is less than the mean life of a new item.

We need only show  $E[\hat{\rho}_{r,n}] \leq E[\hat{\rho}_{r,n}^{**}]$ . Let

$$F_u(x) = \frac{F(x+u) - F(u)}{F(u)} \quad \text{and} \quad G(u) = P[X_1 \leq u]. \quad \text{Then}$$

$$P[X_2 - X_1 \geq \frac{x}{n-1} \mid X_1 = u] = [F_u(\frac{x}{n-1})]^{n-1} \quad \text{and}$$

$$E[(n-1)(X_2 - X_1)] = \int_0^\infty \int_0^\infty [F_u(\frac{x}{n-1})]^{n-1} dx dG(u). \quad \text{Similarly}$$

under Policy ..

$$P[X_2^{**} - X_1^{**} \geq \frac{x}{n} \mid X_1 = u] = [F_u(\frac{x}{n})]^n$$

and

$$E[n(X_2^{**} - X_1^{**})] = \int_0^\infty \int_0^\infty [F_u(\frac{x}{n})]^n dx dG(u).$$

Since  $[F_u(\frac{x}{n})]^n \geq [F_u(\frac{x}{n-1})]^{n-1}$  when  $F$  is IFR we have

$$E[(n-1)(X_2 - X_1)] \leq E[n(X_2^{**} - X_1^{**})].$$

To show

$$E[(n-i+1)(X_i - X_{i-1})] \leq E[(n-i+1)(X_i^{**} - X_{i-1}^{**})]$$

for  $2 \leq i \leq n$  we proceed as above except that the definition of  $G$  is different for the two policies. For example, for  $i = 3$  let

$$\begin{aligned} G(u_2 \mid u_1) &= P[X_2 \leq u_2 \mid X_1 = u_1] \\ &= 1 - [F_{u_1}(u_2 - u_1)]^{n-1} \end{aligned}$$

$$\begin{aligned} \text{and} \quad G^*(u_2 \mid u_1) &= P[X_2^{**} \leq u_2 \mid X_1^{**} = u_1] \\ &= 1 - [F_{u_1}(u_2 - u_1)]^n. \end{aligned}$$

Since  $G(u_2 \mid u_1) \leq G^*(u_2 \mid u_1)$  we have

$$\begin{aligned}
& E[(n-2)(X_3 - X_2)] \\
&= \int_0^\infty \int_0^\infty |\bar{F}_{u_2}(\frac{x}{n-2})|^{n-2} dx dG(u_2 | u_1) dG(u_1) \\
&\leq \int_0^\infty \int_0^\infty |\bar{F}_{u_2}(\frac{x}{n-2})|^{n-2} dx dG^*(u_2 | u_1) dG(u_1) \\
&\leq \int_0^\infty \int_0^\infty |\bar{F}_{u_2}(\frac{x}{n})|^n dx dG^*(u_2 | u_1) dG(u_1) \\
&= E[n(X_3^* - X_2^*)] . \quad \text{A similar conditioning argument}
\end{aligned}$$

works for  $3 \leq i \leq n-1$  .  $\square$

Bounds on estimates for the reliability function. The minimum variance unbiased estimate for  $R(t) = \bar{F}(t)$  ( $t$  is fixed) under the exponential assumption is

$$\hat{R}_1(t) = \max[0, (1 - \frac{t}{Z_r})^{r-1}]$$

where  $Z_r = \sum_{i=1}^r X_i + (n-r)X_r$  . For a discussion of such minimum variance unbiased estimates see Tate (1959). For convenience, assume that  $\theta = 1$  . Then, under the exponential assumption,  $Z_r$  has density

$$g_r(y) = \frac{y^{r-1} e^{-y}}{(r-1)!} .$$

Theorem 2.6. If  $F$  is IFR with mean  $\theta = 1$  and  $t < \theta = 1$ , then

$$E[\hat{R}_1(t)] \geq \int_t^1 [1 - \frac{t}{y}]^{r-1} g_r(y) dy + \int_1^\infty (1-t)^{r-1} g_r(y) dy .$$

Proof. Without loss of generality we may assume  $F$  continuous (see preliminaries). Let  $\phi^{-1}(y) = -\log \bar{F}(y)$  . Then  $\phi$  is concave, increasing and  $\phi(0) = 0$  . If  $Y_i$  is the  $i^{\text{th}}$  order statistic

from an exponentially distributed random variable with  $\theta = 1$ , then  $X_i = \phi(Y_i)$  is the  $i^{\text{th}}$  order statistic from an IFR random variable with distribution  $F$  and mean  $\theta = 1$ . Furthermore

$$\begin{aligned} Z_r &= \sum_{i=1}^r \phi(Y_i) + (n-r) \phi(Y_r) \\ &\geq \phi \left[ \sum_{i=1}^r Y_i + (n-r) Y_r \right] = \phi [Z_r^*] \end{aligned}$$

by the previous lemma. Therefore

$$\begin{aligned} \hat{R}_1(t) &= \max \left[ 0, \left( 1 - \frac{t}{Z_r} \right)^{r-1} \right] \\ &\geq \max \left[ 0, \left( 1 - \frac{t}{\phi(Z_r^*)} \right)^{r-1} \right]. \end{aligned}$$

Since

$$\phi(y) \geq \begin{cases} y & y < 1 \\ 1 & y \geq 1 \end{cases}$$

we have

$$E[\hat{R}_1(t)] \geq \int_t^1 \left[ 1 - \frac{t}{y} \right]^{r-1} g_r(y) dy + \int_1^\infty (1-t)^{r-1} g_r(y) dy \quad (2.1)$$

The maximum likelihood estimate for  $R(t)$  under the exponential assumption is

$$\hat{R}_2(t) = e^{-\frac{t}{\hat{\theta}_{r,n}}}$$

where  $\hat{\theta}_{r,n}$  was defined in (2.1). Pugh (1963) has shown that under the exponential assumption  $\hat{R}_2(t)$  is negatively biased when the true reliability  $R(t) > \frac{1}{e} \approx .368$ . Assuming  $F$  is IFR we can obtain a lower bound on  $E[\hat{R}_2(t)]$ .

Theorem 2.7. If  $F$  is IFR with mean  $\theta = 1$ , then



$$E[\hat{R}_2(t)] \geq \int_0^1 e^{-\frac{tr}{y\theta}} g_r(y) dy + e^{-\frac{tr}{\theta}} \int_1^{\infty} g_r(y) dy.$$

The proof parallels that of Theorem 2.6.

Estimates and confidence bounds on percentiles. If  $F$  is IFR with mean  $\theta$  and  $p^{\text{th}}$  percentile  $\xi_p$ , then

$$[-\log(1-p)]\theta \leq \xi_p \leq \frac{[-\log(1-p)]}{p} \theta.$$

(See Barlow and Marshall (1964)). Hence by Theorem 2.1

$$E\left[\hat{\theta}_{r,n} \frac{[-\log(1-p)]}{p}\right] \geq \xi_p \quad \text{for } r = 1, 2, \dots, n$$

while

$$E\left[\hat{\theta}_{n,n} [-\log(1-p)]\right] \leq \xi_p$$

and one might be tempted to use these estimates to bracket  $\xi_p$ .

Intuitively, we want a confidence interval to have small expected width when it covers the true percentile. The usual distribution-free confidence intervals based on order statistics have smaller conditional expected width under the IFR (DFR) assumption than under the exponential assumption, given that the interval contains the true percentile. To see this let  $Y = -\frac{\xi_p \log F(X)}{-\log(1-p)}$  and note that  $Y$  is exponentially distributed with  $p^{\text{th}}$  percentile  $\xi_p$  when  $F$  is continuous. Suppose that  $X_i \leq \xi_p \leq X_j$ . Then clearly

$$\frac{Y_j - Y_i}{X_j - X_i} \geq 1, \quad \text{which implies}$$

$$E[X_j - X_i \mid X_i \leq \xi_p \leq X_j] \leq E[Y_j - Y_i \mid Y_i \leq \xi_p \leq Y_j].$$

3. Estimates based on truncated samples. If  $n$  items are placed on life test and if sampling is terminated at time  $T$ , the associated sample is called a truncated sample. Let  $X_1 \leq X_2 \leq \dots \leq X_n$  denote an ordered sample from a distribution  $F$  and let

$$V(T) = \sum_{i=1}^r X_i + (n - r)T$$

where  $r$  is a random variable and denotes the number of  $X$ 's less than  $T$ . Then  $V(T)$  is the total life observed up to time  $T$ . This statistic occurs, for example, in sequential life tests for the exponential case (Epstein and Sobel, 1955). It is not surprising that this statistic also has greater expected value under the IFR assumption.

For convenience, let  $G(x) = 1 - e^{-\frac{x}{\theta}}$ .

Theorem 3.1. If  $F$  is IFR (DFR) with mean  $\theta$ , then

$$E_F[V(T)] \underset{(<)}{\geq} E_G[V(T)]$$

Proof. Assume  $F$  IFR and let  $X_1 \leq X_2 \leq \dots \leq X_n$  denote an ordered sample from  $F$ . Without loss of generality we may assume  $F$  continuous. Let  $y = -\theta \log \bar{F}(x)$ . We know there exists  $x_0 > \theta$  such that  $x \geq -\theta \log \bar{F}(x)$  for  $x \leq x_0$  and  $x \leq -\theta \log \bar{F}(x)$  for  $x > x_0$  (Barlow and Marshall, 1964).

Let  $Y_i = -\theta \log \bar{F}(X_i)$ . If  $T \leq x_0$ , then

$$\sum_{i=1}^r X_i + (n - r)T \geq \sum_{i=1}^r Y_i + (n - r)T \geq \sum_{i=1}^s Y_i + (n - s)T$$

where  $r$  ( $s$ ) denotes the number of  $X$ 's ( $Y$ 's) less than  $T$ .

Hence for  $T \leq x_0$

$$E_F[V(T)] \geq E_G[V(T)] .$$

Let

$$Y_i^* = \begin{cases} Y_i & \text{if } Y_i \leq T \\ T & \text{otherwise.} \end{cases}$$

For  $T > x_0$ ,

$$\begin{aligned} & \sum_{i=1}^r X_i + (n-r)T - \sum_{i=1}^s Y_i - (n-s)T \\ &= \sum_{i=1}^r X_i + (n-r)T - \sum_{i=1}^r Y_i^* - (n-r)T \\ &\geq \sum_{i=1}^r (X_i - Y_i) + \sum_{i=r+1}^n (X_i - Y_i) \end{aligned}$$

since  $X_i \leq Y_i$  for  $i > r$ . Hence

$$E_F[V(T)] - E_G[V(T)] \geq E\left[\sum_{i=1}^n X_i\right] - E\left[\sum_{i=1}^n Y_i\right] = 0$$

for  $T > x_0$ .

A similar argument holds for the DFR case.

Consider the estimate

$$\hat{\theta}(T) = \frac{V(T)}{r} = \begin{cases} \frac{\sum_{i=1}^r X_i + (n-r)T}{r} & \text{if } r > 0 \\ nT & \text{if } r = 0 \end{cases}$$

When  $F$  is the exponential distribution,  $\hat{\theta}(T)$  is the maximum likelihood estimate of  $\theta$ . In this case

$$E[\hat{\theta}(T)] = \theta - \frac{\text{cov}(r, \hat{\theta}(T))}{1 - \exp(-\frac{T}{\theta})} > \theta$$

since  $r$  and  $\hat{\theta}(T)$  are negatively correlated (Bartholomew, 1957). In the IFR case, this statistic exhibits even greater bias for  $T < \theta$ . As before let  $G(x) = 1 - e^{-x/\theta}$ .

Theorem 3.2. If  $F$  is IFR with mean  $\theta$ , but not degenerate then

$$E_F[\hat{\theta}(T) \mid r \geq 1] \geq E_G[\hat{\theta}(T) \mid r \geq 1] \quad \text{for } T < \theta.$$

Proof. Assume  $F$  is continuous and let  $Y_i = -\theta \log \bar{F}(X_i)$  as

before and let

$$a_i = \begin{cases} 1 & \text{if } X_i \leq T \\ 0 & \text{otherwise} \end{cases}$$

$$b_i = \begin{cases} 1 & \text{if } Y_i \leq T \\ 0 & \text{otherwise} \end{cases}.$$

We can write

$$\hat{\theta}(T) = \frac{\sum_{i=1}^n [T - a_i(T - X_i)]}{\sum_{i=1}^n a_i} \quad \text{if } \sum_{i=1}^n a_i \geq 1.$$

Assume  $T < \theta$ . As in the previous proof  $X_i < T$  implies  $Y_i < T$  and hence  $a_i \leq b_i$ . If  $a_i = 1$ , then  $b_i = 1$  and

$T - X_i < T - Y_i$ . Hence if  $\sum_{i=1}^n a_i \geq 1$ ,

$$\frac{\sum_{i=1}^n [T - a_i(T - X_i)]}{\sum_{i=1}^n a_i} \geq \frac{\sum_{i=1}^n [T - b_i(T - Y_i)]}{\sum_{i=1}^n b_i}$$

and

$$E_F[\hat{\theta}(T) \mid r \geq 1] \geq E_G[\hat{\theta}(T) \mid r \geq 1] \quad \text{for } T < \theta. \parallel$$

### Inverse Binomial Sampling.

Nadler (1960) has considered the following type of sampling.

An item having life distribution  $F$  with mean  $\theta$  is put on test until it fails or time  $t$  has elapsed; at this time the item is replaced by a fresh item. This is repeated sequentially until  $r$  actual failures are observed. The number  $N$  of items that have to be tested until the  $r$  actual failures are obtained is a random variable. Nadler

(1960) showed that when  $F(x) = 1 - e^{-\frac{x}{\theta}}$ , an unbiased estimate of  $\theta$  is

$$\hat{\theta}_r(t) = \frac{1}{r} \left[ \sum_{i=1}^r Y_i + (N-r)t \right]$$

where the  $Y_1, \dots, Y_r$  are the  $r$  life lengths not exceeding  $t$ .

We show next that when  $F$  is IFR (DFR) with mean  $\theta$ , then  $\hat{\theta}_r(t)$  is biased high (low).

**Theorem 3.3.** If  $F$  is IFR (DFR) with mean  $\theta$ , then  $E\hat{\theta}_r(t) \geq (\leq) \theta$ .

**Proof.** Let  $F$  be IFR. Let  $Z_i$  denote test time elapsed between the  $i-1^{\text{st}}$  failure time and the  $i^{\text{th}}$  failure time,  $i = 1, 2, \dots, r$ , where the  $0^{\text{th}}$  failure time is defined to be 0. Then

$$\hat{\theta}_r(t) = \frac{1}{r} \sum_{i=1}^r Z_i.$$

Next consider an alternate testing procedure differing in that replacement occurs only upon failure. Let  $Z'_i$  = test time elapsed between the  $i-1^{\text{st}}$  failure and the  $i^{\text{th}}$  failure under the alternate testing procedure. Now since  $F$  is IFR,  $Z_i$  is stochastically larger than  $Z'_i$ . It follows that

$$E\hat{\theta}_r(t) = \frac{1}{r} \sum_{i=1}^r E Z_i \geq \frac{1}{r} \sum_{i=1}^r E Z_i' = \theta$$

The inequality is reversed when  $F$  is DFR.

Sampling with replacement. In this case

$$\hat{\theta}(T) = \frac{nT}{r} = \frac{nT}{\sum_{i=1}^n N_i(T)}$$

where  $N_i(T)$  denotes the number of replacements in the  $i^{\text{th}}$  item position and  $r = \sum_{i=1}^n N_i(T)$  denotes the total number of replacements in  $[0, T]$ . Of course  $E\left[\frac{1}{\sum_{i=1}^n N_i(T)}\right]$  is unbounded.

However, we know that  $E[N_i(T)] \leq \frac{T}{\theta}$  for all  $T > 0$  (Barlow and Proschan, 1964b). Hence this again indicates that  $\hat{\theta}(T)$  will tend to be larger in the IFR case than in the exponential case.

4. Bounds on time to  $r^{\text{th}}$  failure. Under the exponential assumption, the distribution of the statistic  $\hat{\theta}_{r,n}$  depends only on  $r$  and not on  $n$ . The choice of  $n$  in this case is usually determined by the ratio

$$\frac{E(X_{r,n})}{E(X_{r,r})}$$

which is an indirect measure of the expected saving in time due to putting more than  $r$  items on test but terminating at the  $r^{\text{th}}$  failure (Epstein, 1960a). We always have

$$\frac{E(X_{r,n})}{E(X_{r,r})} \leq 1.$$

Since the bound is attained by the degenerate distribution (which is the limit of IFR distributions), this is not a useful measure if we assume only IFR. However, we can obtain non-trivial bounds on  $E(X_{r,n})$ .

Assume  $F$  is IFR, with mean 1, and continuous. We may write  $Y_i = -\log \bar{F}(X_i)$  where  $Y_i$  is the  $i^{\text{th}}$  order statistic in a sample of  $n$  from distribution  $G(x) = 1 - e^{-x}$ , and is a convex function of  $X_i$ , where  $X_i$  is the  $i^{\text{th}}$  order statistic in a sample of  $n$  from  $F$ . By Jensen's inequality

$$E(Y_i) \geq -\log \bar{F}[E(X_i)],$$

so that

$$\bar{F}[E(X_i)] \geq e^{-E(Y_i)}.$$

If  $b(x)$  is a sharp upper bound on  $\bar{F}(x)$ , then  $b$  is decreasing to 0 and

$$b[E(X_i)] \geq e^{-E(Y_i)}.$$

Hence choosing  $x_0$  such that

$$b(x_0) = e^{-E(Y_i)},$$

where of course

$$E(Y_i) = \sum_{j=1}^i \frac{1}{(n-j+1)},$$

we have

$$E(X_i) \leq x_0.$$

Using tabled upper bounds on  $\bar{F}$  given one or two moments of  $F$  (Barlow and Marshall, 1963) we can obtain upper bounds on  $E(X_i)$ .

When  $F$  is DFR we can, in a similar manner, obtain lower bounds on  $E(X_i)$  using lower bounds on  $\bar{F}$ .

If we specify the first moment of  $F$ , explicit upper bounds can be given on  $E(X_i)$  when  $F$  is IFR, as shown in

Theorem 4.1. If  $X_1 \leq X_2 \leq \dots \leq X_n$  are order statistics from an IFR random variable with mean  $\theta$  and  $Y_1 \leq Y_2 \leq \dots \leq Y_n$  are the order statistics from  $G(x) = 1 - e^{-x}$ , then

$$(a) \quad \theta E(Y_1) \leq E(X_1) \leq \theta$$

$$(b) \quad E(X_i) \leq \frac{\theta E(Y_i)}{1 - e^{-E(Y_i)}} \quad 1 < i < n$$

$$(c) \quad \theta \leq F(X_n) \leq \theta E(Y_n)$$

(a) and (c) are sharp and (b) is non-trivial though not sharp.

Proof. (a) and (c) are shown in Barlow and Proschan (1964a),

Chapter 2. Hence we need only prove (b) and we may assume  $\theta = 1$ .

First let us verify that (b) is non-trivial. Note that by (a),  $EX_{n-1} \leq EX_n \leq EY_n$ , so that  $EY_n$  is a trivial upper bound for  $EX_{n-1}$ . Therefore, a non-trivial upper bound for  $EX_{n-1}$  must be

less than  $EY_n$ ; i.e., we must show that  $\frac{EY_{n-1}}{1 - e^{-EY_{n-1}}} < EY_n$ .

But for  $z > 0$ ,  $\frac{z}{1 - e^{-z}} < z + 1$ ; thus letting  $z = EY_{n-1}$

$\frac{1}{n} + \frac{1}{n-1} + \dots + \frac{1}{2}$ , we conclude

$$\frac{EY_{n-1}}{1 - e^{-EY_{n-1}}} < \frac{1}{n} + \frac{1}{n-1} + \dots + \frac{1}{2} + 1 = EY_n.$$

To show (b) use the bound

$$F(x) \leq b(x) = \begin{cases} 1 & x \leq \theta \\ e^{-wx} & x > \theta \end{cases}$$



where  $w$  depends on  $x$  and satisfies  $\int_0^x e^{-wu} du = \theta$ .  
(Barlow and Marshall, 1964).||

Sharp bounds will be derived in a future publication. However these are not as convenient as the bounds of Theorem 4.1.

Bounds on expected values of order statistics can also be given in terms of the  $p^{\text{th}}$  percentile.

Theorem 4.2. Let  $X_1 \leq X_2 \leq \dots \leq X_n$  denote the order statistics from  $F$ , IFR with  $p^{\text{th}}$  percentile  $\xi_p$ . Then

$$(4.1) \quad E(X_j) \leq \max \left\{ \xi_p, \frac{\xi_p}{-\log q} \left( \frac{1}{n} + \dots + \frac{1}{(n-j+1)} \right) \right\}$$

and

$$(4.2) \quad E(X_j) \geq \sum_{i=0}^{j-1} \binom{n}{i} \int_0^{\xi_p} \left( 1 - e^{-\frac{x \log q}{\xi_p}} \right)^i \left( e^{-\frac{x \log q}{\xi_p}} \right)^{n-i} dx$$

where  $q = 1 - p$ . All inequalities are sharp.

Proof. To show (4.1), let

$$\bar{G}_\Delta(x) = \begin{cases} 1 & 0 \leq x \leq \Delta \\ q \exp \left[ \left( \frac{x - \xi_p}{\xi_p - \Delta} \right) \log q \right] & x > \Delta \end{cases}$$

Note that  $\bar{G}_\Delta(\Delta) = 1$  and  $\bar{G}_\Delta(\xi_p) = 1 - p = q$ . Since  $\log \bar{F}(x)$  is concave, there exists at least one value of  $\Delta \geq 0$  such that  $\bar{G}_\Delta(x) \geq \bar{F}(x)$  for all  $x \geq 0$ . Thus  $E(X_j) \leq \sup_{0 \leq \Delta \leq \xi_p} E(Y_j)$  where

$Y_j$  is the  $j^{\text{th}}$  order statistic from  $G_\Delta$ . Now

$$\begin{aligned} E(Y_j) &= \Delta + \int_\Delta^\infty \sum_{i=0}^{j-1} \binom{n}{i} [G_\Delta(x)]^i [\bar{G}_\Delta(x)]^{n-i} dx \\ &= \Delta + \int_\Delta^\infty \frac{\Gamma(n+1)}{\Gamma(j)\Gamma(n+1-j)} \int_{G_\Delta(x)}^1 t^{j-1} (1-t)^{n-j} dt dx \end{aligned}$$

To find the maximizing  $\Delta$ , consider

$$\begin{aligned} \frac{\partial}{\partial \Delta} E(Y_j) &= 1 - \frac{\Gamma(n+1)}{\Gamma(j)\Gamma(n+1-j)} \int_{G_\Delta(\Delta)}^1 t^{j-1}(1-t)^{n-j} dt \\ &\quad + \frac{\Gamma(n+1)}{\Gamma(j)\Gamma(n+1-j)} \int_{\Delta}^{\infty} [G(x)]^{j-1} [\bar{G}(x)]^{n-j} \\ &\quad \cdot q \exp\left[\frac{x - \xi_p}{\xi_p - \Delta} \log q\right] \log q \frac{x - \xi_p}{(\xi_p - \Delta)^2} dx. \end{aligned}$$

Since  $G_\Delta(\Delta) = 0$ ,  $-(\xi_p - \Delta) \frac{\partial E(Y_j)}{\partial \Delta}$  reduces to

$$\frac{\Gamma(n+1)}{\Gamma(j)\Gamma(n+1-j)} \int_{\Delta}^{\infty} [G_\Delta(x)]^{j-1} [\bar{G}_\Delta(x)]^{n-j} g_\Delta(x) (x - \xi_p) dx = E(Y_j) - \xi_p$$

where  $g_\Delta$  is the density of  $G_\Delta$ .

Hence

$$\begin{aligned} -(\xi_p - \Delta) \frac{\partial}{\partial \Delta} E(Y_j) &= \Delta - \frac{\xi_p - \Delta}{\log q} \left( \frac{1}{n} + \dots + \frac{1}{n-j+1} \right) - \xi_p \\ &= -(\xi_p - \Delta) \left[ 1 + \frac{1}{\log q} \left( \frac{1}{n} + \dots + \frac{1}{n-j+1} \right) \right]. \end{aligned}$$

For  $j$  such that  $1 + \frac{1}{\log q} \left( \frac{1}{n} + \dots + \frac{1}{n-j+1} \right) \leq 0$ , we have

$$\frac{\partial}{\partial \Delta} E(Y_j) \leq 0. \text{ For } j \text{ such that } 1 + \frac{1}{\log q} \left( \frac{1}{n} + \dots + \frac{1}{n-j+1} \right) \geq 0$$

we have  $\frac{\partial}{\partial \Delta} E(Y_j) \geq 0$ . Thus  $E(Y_j)$  is maximized in the first

case at  $\Delta = 0$  and at  $\Delta = \xi_p$  in the second case. When  $\Delta = 0$ ,

$$E(Y_j) = \frac{\xi_p}{-\log q} \left( \frac{1}{n} + \dots + \frac{1}{n-j+1} \right). \text{ When } \Delta = \xi_p, E(Y_j) = \xi_p.$$

To show (4.2). Let  $\bar{G}(x) = \begin{cases} e^{x \log \frac{q}{\xi_p}} & \text{for } 0 \leq x < \xi_p \\ 0 & \text{for } \xi_p \leq x < \infty. \end{cases}$

Then  $\bar{G}(x) \leq \bar{F}(x)$  for all  $x \geq 0$  and  $G$  has  $p^{\text{th}}$  percentile  $\xi_p$ . Thus  $E(Y_j) \leq E(X_j)$  where  $Y_j$  is the  $j^{\text{th}}$  order statistic from  $G$ . But

$$E(Y_j) = \int_0^\infty \sum_{i=0}^{j-1} \binom{n}{i} [G(x)]^i [\bar{G}(x)]^{n-i} dx.$$

Using the above definition of  $G$ , we obtain (4.2).  $\square$

5. Further results on order statistics and spacings. In this section we present some results of theoretical interest concerning order statistics and their spacings from  $PF_2$  and IFR (DFR) distributions. Many of the results hold without the restriction  $F(0^-) = 0$ .

First we present some total positivity properties of the order statistics. A function  $K(x, y)$  of two real variables is said to be totally positive of order  $r$  ( $TP_r$ ) if for all  $1 \leq m \leq r$ ;  $x_1 \leq x_2 \leq \dots \leq x_m$  and  $y_1 \leq y_2 \leq \dots \leq y_m$  we have the determinant inequalities

$$\left| K(x_i, y_j) \right|_{i,j=1}^m \geq 0.$$

The following lemma is of use later on, as well as of interest in its own right.

Lemma 5.1. Let  $F$  be a distribution having density  $f$  with  $f(x)$  not necessarily 0 for  $x < 0$ . Let  $f_i(x)$  be the density of the  $i^{\text{th}}$  order statistic in a sample of size  $n$ . Then  $f_i(x)$  is  $TP_0$  in  $i, x$  where  $i = 1, 2, \dots, n$  and  $-\infty < x < \infty$ .

Proof.

$$(5.1) \quad f_i(x) = \frac{n!}{(i-1)!(n-i)!} F^{i-1}(x) \bar{F}^{n-i}(x) f(x).$$

Since  $\left\{ \frac{F(x)}{\bar{F}(x)} \right\}^{i-1}$  is  $TP_{\infty}$  in  $i$  and  $x$ , when  $i = 1, 2, \dots, n$  and  $-\infty < x < \infty$ , the conclusion follows. ||

We may obtain a similar result concerning the right hand tail of the distribution of an order statistic.

Lemma 5.2. Let  $F$  be any distribution with  $F(x)$  not necessarily 0 for  $x < 0$ ,  $F_i$  the corresponding distribution of the  $i^{\text{th}}$  order statistic. Then  $\bar{F}_i(x)$  is  $TP_{\infty}$  in  $i, x$ , where  $i = 1, 2, \dots, n$  and  $-\infty < x < \infty$ .

Proof.

$$(5.2) \quad \bar{F}_i(x) = \sum_{j=0}^{i-1} \binom{n}{j} F^j(x) \bar{F}^{n-j}(x).$$

Now  $\left\{ \frac{F(x)}{\bar{F}(x)} \right\}^j$  is  $TP_{\infty}$  in  $x, j$ . Therefore

$$\bar{F}_i(x) = \sum_{j=0}^{\infty} \binom{n}{j} F^n(x) \left\{ \frac{F(x)}{\bar{F}(x)} \right\}^j H(i-1-j) \text{ is also}$$

$TP_{\infty}$  in  $i$  and  $x$ , where  $H(k) = 1$  for  $k \geq 0$ , 0 otherwise. ||

In Barlow, Marshall, Proschan (1963), it is shown that the order statistics from an IFR distribution themselves have an IFR distribution. The next lemma shows a similar preservation of the  $PF_2$  property.

Lemma 5.3. Suppose the underlying density  $f$  is  $PF_2$ , with  $f(x)$  not necessarily 0 for  $x < 0$ . Then the density  $f_i$  of the  $i^{\text{th}}$  order statistic is also  $PF_2$  for fixed  $i = 1, 2, \dots, n$ .

Proof. It is easy to verify that when  $f$  is  $PF_2$ , so is  $F$  and  $\bar{F}$ . Thus  $\log f$ ,  $\log F$ , and  $\log \bar{F}$  are concave. It follows from (5.1) that  $\log f_i$  is concave, or equivalently,  $f_i$  is  $PF_2$  for fixed  $i = 1, 2, \dots, n$ . ||

Next we obtain some comparisons between the order statistics of an IFR (DFR) distribution and the corresponding order statistics of an exponential distribution.

Theorem 5.4. Let  $X_1 \leq X_2 \leq \dots \leq X_n$ ,  $n \geq 2$ , be order statistics from  $F$ , an IFR (DFR) distribution with mean  $\theta$ , but

$\neq 1 - e^{-\frac{t}{\theta}}$ . Let  $Y_1 \leq Y_2 \leq \dots \leq Y_n$  be order statistics from

$G(t) = 1 - e^{-\frac{t}{\theta'}}$ . Then

- (a)  $EX_j - EY_j$  has at most one change of sign as  $j$  goes from 1 to  $n$ . Moreover if one change of sign does occur, then  $EX_j - EY_j$  goes from positive (negative) to negative (positive) values.
- (b) If  $\theta = \theta'$ , then one change of sign does occur.

Proof.

(a) Assume  $F$  is a continuous IFR distribution with mean  $\theta$  and  $G$  is exponential with mean  $\theta'$ . We have seen in Section 1 that if  $Y$  has distribution  $G$ , then  $X = \phi\left(\frac{Y}{\theta'}\right)$  has distribution  $F$ , where  $\phi^{-1}(x) = -\log \bar{F}(x)$ , a convex increasing function which is 0 for  $x = 0$ .

Thus

$$EX_i - EY_i = \int_0^\infty \left\{ \phi\left(\frac{y}{\theta'}\right) - y \right\} g_i(y) dy,$$

where  $g_i$  is the density of the  $i^{\text{th}}$  order statistic from the exponential distribution  $G$ . By Lemma 5.1,  $g_i(y)$  is  $TP_\infty$  in  $y$  and  $v$ . Also  $\phi(\frac{y}{\theta}) - y$  changes sign at most once, and if once, from positive to negative values. By the variation diminishing property of totally positive functions (Karlin, 1964, p. 34),  $EX_i - EY_i$  also changes sign at most once, and from positive to negative values, if at all.

(b) If  $\theta = \theta'$ , then  $\sum_{i=1}^n EX_i = \theta = \theta' = \sum_{i=1}^n EY_i$ . Hence  $EX_i - EY_i$

must change sign at least once or be identically 0 for

$i = 1, 2, \dots, n$ . Now since  $F \not\equiv G$ ,  $F$  cannot agree with  $G$  on an interval. Hence by Corollary 4.10, Chapter 2 of Barlow and Proschan (1964),  $EX_1 > EY_1$  and  $EX_n < EY_n$ . Thus  $EX_i - EY_i$  is not identically 0 for  $i = 1, 2, \dots, n$ . Hence  $EX_i - EY_i$  changes sign exactly once.

If  $F$  is IFR but not continuous, we may obtain the same result by using continuous IFR approximations.

Finally, a similar argument holds if  $F$  is DFR.  $\parallel$

Actually, under the same hypothesis we may prove a stronger version of (a) in which  $EX_i - EY_i$  is replaced by  $EX_i^\alpha - EY_i^\alpha$ ,  $\alpha > 0$ . If instead of assuming  $EX = EY$  in (b), we assume  $EX^\alpha = EY^\alpha$ , then we may show that one change of sign of  $EX_i^\alpha - EY_i^\alpha$  does occur. We omit the details.

We may obtain further consequences of Theorem 5.4 using the notion of majorization. A vector  $\underline{a} = (a_1, a_2, \dots, a_n)$  majorizes a vector  $\underline{b} = (b_1, b_2, \dots, b_n)$  (written  $\underline{a} > \underline{b}$ ) if

$a_1 \geq a_2 \geq \dots \geq a_n, b_1 \geq b_2 \geq \dots \geq b_n, \sum_{i=1}^k a_i \geq \sum_{i=1}^k b_i,$   
 $k = 1, 2, \dots, n-1, \text{ and } \sum_{i=1}^n a_i = \sum_{i=1}^n b_i.$  See Hardy, Littlewood,  
 Pólya (1952), Chapter II.

Theorem 5.5. Let  $X_1 \leq \dots \leq X_n$  be order statistics from  $F$ ,  
 an IFR (DFR) distribution with mean  $\theta$ ,  $Y_1 \leq \dots \leq Y_n$  be  
 order statistics from  $G(t) = 1 - e^{-\frac{t}{\theta}}$ . Then  $(EY_n, EY_{n-1}, \dots, EY_1)$   
 $>(<)(EX_n, EX_{n-1}, \dots, EX_1)$ .

Proof. Let  $F$  be IFR. From Theorem 5.4 we know

$EY_{n-i+1} - EX_{n-i+1}$  has one change of sign, from plus to minus as  
 $i$  goes from 1 to  $n$ . We also know  $\sum_{i=1}^n EY_{n-i+1} = n\theta = \sum_{i=1}^n EX_{n-i+1}$ .

Thus  $\sum_{i=1}^j EY_{n-i+1} \geq \sum_{i=1}^j EX_{n-i+1}, j = 1, 2, \dots, n$ . Finally,  $EY_{n-i+1}$

and  $EX_{n-i+1}$  are decreasing in  $i$ . Thus the conclusion follows.

A similar argument holds if  $F$  is DFR. ||

Using Karamata's Theorem we obtain Theorem 5.6 below.

Karamata's Theorem states that if  $\psi$  is continuous and convex and

$\underline{a} > \underline{b}$ , then

$$(5.3) \quad \sum_{i=1}^n \psi(a_i) \geq \sum_{i=1}^n \psi(b_i).$$

See Hardy, Littlewood, Pólya (1952), p. 89.

Theorem 5.6. Let  $\psi$  be continuous and convex,  $X_1, \dots, X_n$ ,

$Y_1, \dots, Y_n$  as in Theorem 5.5. Then

$$\sum_{i=1}^n \psi(EY_i) \geq \sum_{i=1}^n \psi(EX_i).$$

Proof. Let  $F$  be IFR. By Theorem 5.5,  $(FY_n, \dots, FY_1) > (EX_n, \dots, EX_1)$ . Hence by Karamata's Theorem, the conclusion follows. A similar argument holds if  $F$  is DFR.  $\square$

Using Theorem 5.5 we obtain

Theorem 5.7. Let  $c_1 \leq c_2 \leq \dots \leq c_n$ ,  $X_1, \dots, X_n$ ,  $Y_1, \dots, Y_n$  as in Theorem 5.5. Then  $\sum_{i=1}^n c_i FY_{n-i+1} \geq \sum_{i=1}^n c_i EX_{n-i+1}$ .

Proof. Let  $F$  be IFR. Defining  $d_i = FY_{n-i+1} - EX_{n-i+1}$ , write

$$\begin{aligned} \sum_{i=1}^n c_{n-i+1} (FY_{n-i+1} - EX_{n-i+1}) &= (c_n - c_{n-1})d_1 + (c_{n-1} - c_{n-2})(d_1 + d_2) \\ &+ (c_{n-2} - c_{n-3})(d_1 + d_2 + d_3) + \dots + (c_2 - c_1)(d_1 + \dots + d_{n-1}) \\ &+ c_1(d_1 + \dots + d_n). \end{aligned}$$

Since  $c_i - c_{i+1} \geq 0$ ,  $i = 1, \dots, n-1$ ,  $d_1 + \dots + d_j \geq 0$ ,  $j = 1, \dots, n-1$ , and  $d_1 + \dots + d_n = 0$ , we conclude that

$$\sum_{i=1}^n c_{n-i+1} (FY_{n-i+1} - EX_{n-i+1}) \geq 0.$$

A similar argument holds when  $F$  is DFR.  $\square$

Finally we summarize some results concerning the covariance of order statistics obtained by Tukey (1958). He shows that if  $F$  is IFR, then for  $h \leq j < k$ ,  $\text{cov}(X_k, X_h) \leq \text{cov}(X_j, X_h)$ , where  $X_j$  is the  $j^{\text{th}}$  order statistic from  $F$ . He further shows that if  $F$  satisfies both

- (a)  $\log \bar{F}$  is concave (i.e.,  $F$  is IFR), and
- (b)  $\log F$  is concave,



then

(1) the covariance of any two order statistics is less than the variance of either, and

(2) the covariance between order statistics  $X_j, X_k$  is monotone in  $j$  and  $k$  separately, decreasing as  $j$  and  $k$  separate from one another.

Note that if  $f$  is  $PF_2$ , then  $F$  satisfies both (a) and (b) above.

Next we derive properties of the spacings  $X_1, X_2 - X_1, \dots, X_n - X_{n-1}$  from  $PF_2$  and IFR (DFR) distributions similar to those of the order statistics  $X_1 \leq X_2 \leq \dots \leq X_n$  obtained above. We first consider total positivity properties.

Theorem 5.8. Let  $f$  be  $PF_2$  with  $f(x)$  not necessarily 0 for  $x < 0$ . Then  $h_i$ , the density of  $X_i - X_{i-1}$ , is  $PF_2$  for fixed  $i = 2, 3, \dots, n$ . If we assume further that  $f(x) = 0$  for  $x < 0$ , then  $h_1$  is  $PF_2$ , where  $h_1$  is the density of  $X_1$ .

Proof.

$$(5.4) \quad h_i(x) = \frac{n!}{(i-2)!(n-i)!} \int F^{i-2}(u) f(u) f(u+x) \bar{F}^{n-i}(u+x) du$$

for  $i = 2, 3, \dots, n$ .

Since  $f$  is  $PF_2$ , so is  $r(u) = F^{i-2}(-u)f(-u)$ ,  $s(u) = f(u)\bar{F}^{n-i+1}(u)$ . Hence so is

$$h_i(x) = \frac{n!}{(i-1)!(n-i+1)!} \int r(-u)s(u+x) du$$

for fixed  $i = 2, 3, \dots, n$ .

Assuming  $f(x) = 0$  for  $x < 0$ , we see that  $h_1$  is  $PF_2$ , from

$$(5.5) \quad h_1(x) = n f(x) \bar{F}^{n-1}(x) \quad . \parallel$$

Theorem 5.9. If  $F$  is DFR, then  $H_i$  is DFR for fixed  $i = 1, 2, \dots, n$ .

Proof. Since DFR is preserved under convex combinations (Barlow, Marshall, Proschan, 1963, p. 381) we see from the representation

$$(5.6) \quad H_i(x) = \frac{n!}{(i-1)!(n-i)!} \int_0^\infty F^{i-1}(u) f(u) \bar{F}_u^{n-i}(x) du,$$

where  $\bar{F}_u(x) = F(u+x)$ , DFR in  $x$  for fixed  $u$ , that  $H_i$  is DFR for fixed  $i = 1, 2, \dots, n$ . ||

Theorem 5.10. Let  $F$  be DFR with  $F(x) < 1$  for all  $x > 0$ . Then  $H_i(x)$  is  $TP_2$  in  $i, x$  where  $i = 2, 3, \dots, n$  and  $x \geq 0$ .

Proof.  $F^{i-2}(u) f(u)$  is  $TP_2$  in  $i = 2, 3, \dots, n$  and  $u \geq 0$ .

$\bar{F}^{n-i+1}(u+x)$  is  $TP_2$  in  $i, u$ , in  $i, x$ , and in  $u, x$  ( $u \geq 0, x \geq 0$ ).

Thus by a theorem\* in the book by Karlin (in process)

$$(5.6) \quad H_i(x) = \frac{n!}{(i-1)!(n-i)!} \int_0^\infty F^{i-2}(u) f(u) \bar{F}^{n-i+1}(u+x) du$$

is  $TP_2$  in  $i, x$ , where  $i = 2, 3, \dots, n$  and  $x \geq 0$ . ||

\* Theorem (Karlin) let  $\lambda, x, \xi$  traverse linear sets  $\Lambda, X, \Xi$  respectively. Suppose  $h(\lambda, x) = \int f(\lambda, x, \xi) g(\lambda, \xi) d\mu(\xi)$  is well defined on  $\Lambda \times X$ , where  $\mu$  is a  $\sigma$ -finite measure, and

(i)  $f(\lambda, x, \xi) > 0$  for all  $\lambda$  in  $\Lambda$ ,  $x$  in  $X$ , and  $\xi$  in  $\Xi$ , and  $f$  is  $TP_2$  for each pair of variables when the third variable is held fixed.

(ii)  $g(\lambda, \xi)$  is  $TP_2$ .

Then  $h$  is  $TP_2$ .

Next we present some majorization properties of the normalized spacings  $(n - i + 1)(X_i - X_{i-1})$ ,  $i = 1, 2, \dots, n$ , similar to those developed above for the order statistics.

Theorem 5.11. Let  $X_1 \leq X_2 \leq \dots \leq X_n$  be the order statistics from  $F$ , an IFR distribution with mean  $\theta$ ,  $Y_1 \leq Y_2 \leq \dots \leq Y_n$  the order statistics from  $G(t) = 1 - e^{-t/\theta}$ . Then

$$\begin{aligned} & E n X_1, E(n-1)(X_2 - X_1), \dots, E(X_n - X_{n-1}) \\ & \succ(\prec) E n Y_1, E(n-1)(Y_2 - Y_1), \dots, E(Y_n - Y_{n-1}) . \end{aligned}$$

Proof. Let  $F$  be IFR. By Theorem 2.1,  $E(n-i+1)(X_i - X_{i-1})$  is decreasing in  $i$ . It is also easy to verify that  $E n Y_1 = E(n-1)(Y_2 - Y_1) = \dots = E(Y_n - Y_{n-1})$ . Since

$$\sum_{i=1}^n E(n-i+1)(X_i - X_{i-1}) = \theta = \sum_{i=1}^n E(n-i+1)(Y_i - Y_{i-1}) ,$$

it follows that

$$\sum_{i=1}^j E(n-i+1)(X_i - X_{i-1}) \geq \sum_{i=1}^j E(n-i+1)(Y_i - Y_{i-1})$$

for  $j = 1, 2, \dots, n-1$ . Thus the conclusion follows.

A similar argument holds if  $F$  is DFR.  $\square$

For normalized spacings, the analogue of Theorem 5.7 is

Theorem 5.12. Let  $c_1 \geq \dots \geq c_n$ ,  $X_1, X_2, \dots, X_n$ ,  $Y_1, Y_2, \dots, Y_n$  as in Theorem 5.11. Then

$$\sum_{i=1}^n c_i E(n-i+1)(X_i - X_{i-1}) \geq \sum_{i=1}^n c_i E(n-i+1)(Y_i - Y_{i-1}) .$$

( $\prec$ )

Proof. The proof parallels that of Theorem 5.7. ||

We immediately obtain:

Corollary 5.13. Let  $X_1, \dots, X_n, Y_1, \dots, Y_n$  be as in Theorem 5.11.

Then for  $1 \leq r \leq n$

$$\sum_{i=1}^r E(n-i+1)(X_i - X_{i-1}) \geq \sum_{i=1}^r E(n-i+1)(Y_i - Y_{i-1}) .$$

Proof. Choose  $c_1 = c_2 = \dots = c_r = 1, c_{r+1} = c_{r+2} = \dots = c_n = 0$  ,

so that  $c_1 \geq c_2 \geq \dots \geq c_n$  . The result follows from Theorem 5.11. ||

## REFERENCES

- Barlow, R. E. and A. Marshall, (1964), "Bounds for Distributions with Monotone Hazard Rate I," Ann. Math. Statist., Vol. 35, No. 3, pp. 1234-1257.
- , (1963), Tables of Bounds for Distributions with Monotone Hazard Rate, Boeing Scientific Research Labs. Doc. DI-82-0249.
- Barlow, R. E., A. W. Marshall, and F. Proschan, (1963), "Properties of Probability Distributions with Monotone Hazard Rate," Ann. Math. Statist., Vol. 34, pp. 375-389.
- Barlow, R. E., and F. Proschan, (1964a), Mathematical Theory of Reliability, J. Wiley and Sons (to appear).
- , (1964b), "Comparison of Replacement Policies and Renewal Theory Implications," Ann. Math. Statist., Vol. 35, No. 2, pp. 577-589.
- Bartholomew, D. J., (1957), "A Problem in Life Testing," J. Amer. Statist. Assn., Vol. 52, pp. 350-355.
- Epstein, B., (1960a), "Statistical Life Test Acceptance Procedures," Technometrics, Vol. 2, No. 4, pp. 435-446.
- , (1960b), "Estimation from Life Test Data," Technometrics, Vol. 2, No. 4, pp. 447-454.
- Epstein, B. and M. Sobel, (1953), "Life Testing," J. Amer. Statist. Assn., Vol. 48, pp. 486-502.
- , (1955), "Sequential Life Tests in the Exponential Case," Ann. Math. Statist., Vol. 26, No. 1, pp. 82-93.
- Hardy, G. H., J. E. Littlewood, and G. Pólya, (1952), Inequalities, 2nd ed., Cambridge University Press.
- Karlin, S., (1964), "Total Positivity, Absorption Probabilities and Applications," Transactions of the American Math. Soc., Vol. 111, pp. 33-107.
- , Total Positivity and Applications to Probability and Statistics, Stanford University Press, Stanford, California, (to appear).

- Marshall, A. W. and F. Proschan, (1964), Maximum Likelihood Estimation for Distributions with Monotone Failure Rate, Boeing Scientific Research Labs. Report DI-82-0329.
- Nadler, J., (1960), "Inverse Binomial Sampling Plans When an Exponential Distribution is Sampled with Censoring," Ann. Math. Statist., Vol. 31, pp. 1201-1204.
- Ostrowski, A., (1952), "Sur Quelques Applications des Fonctions Convexes et Concaves au Sens de I. Schur," J. Math. Pure Appl. Vol. 31, pp. 253-292.
- Proschan, F. and R. Pyke, (1964), Tests for Increasing Failure Rate, Boeing Scientific Research Labs. Doc. (to appear).
- Pugh, E. L., (1963), "The Best Estimate of Reliability in the Exponential Case," Operations Research, No. 1, pp. 57-61.
- Sampling Procedures and Tables for Life and Reliability Testing, (1960), Quality Control and Reliability Handbook (Interim), H108, Office of the Assistant Sec. of Defense, Government Printing Office.
- Tate, R. F., (1959), "Unbiased Estimation Functions of Location and Scale Parameters," Ann. Math. Statist., Vol. 30, No. 2, pp. 362-366.
- Tukey, J. W., (1958), "A Problem of Berkson, and Minimum Variance Orderly Estimators," Ann. Math. Statist., Vol. 29, pp. 588-592.
- Zelen, M., and M. C. Dannemiller, (1961), "The Robustness of Life Testing Procedures Derived from the Exponential Distribution," Technometrics, Vol. 3, pp. 29-49.

**BASIC DISTRIBUTION LIST FOR UNCLASSIFIED  
TECHNICAL REPORTS**

Logistics and Mathematical Statistics Branch Office of Naval Research Washington, D. C. 20360	3 copies
Commanding Officer Office of Naval Research Branch Office Navy #100 Fleet Post Office New York, New York	2 copies
Defense Documentation Center Cameron Station Alexandria, Virginia 22314	20 copies
Defense Logistics Studies Information Exchange Army Logistics Management Center Fort Lee, Virginia Attn: William B. Whichard	1 copy
Technical Information Officer Naval Research Laboratory Washington, D. C. 20390	6 copies
Commanding Officer Office of Naval Research Branch Office 207 West 24th Street New York, New York 10011 Attn: J. Laderman	1 copy
Commanding Officer Office of Naval Research Branch Office 1030 East Green Street Pasadena, California 91101 Attn: Dr. A. R. Laufer	1 copy
Bureau of Supplies and Accounts Code OW Department of the Navy Washington, D. C. 20360	1 copy
Institute for Defense Analyses Communications Research Division von Neumann Hall Princeton, New Jersey	1 copy
Commanding Officer Office of Naval Research Branch Office 495 Summer Street Boston, Massachusetts 02110	1 copy

Commanding Officer  
Office of Naval Research Branch Office  
230 North Michigan Avenue  
Chicago, Illinois 60601

1 copy

Commanding Officer  
Office of Naval Research Branch Office  
1000 Geary Street  
San Francisco, California 94109

1 copy

COR Arthur J. Coyle  
Office of Naval Research - Code 429  
Dept. of the Navy  
Washington, D.C. 20360

1 copy

Mr. William C. Karl  
Office of Naval Research - Code 429  
Dept. of the Navy  
Washington, D.C. 20360

1 copy



University of Chicago  
Statistical Research Center  
Chicago, Illinois 60637  
Attn: Prof. Paul Meier

Stanford University  
Department of Statistics  
Stanford, California 94305  
Attn: Prof. Gerald J. Lieberman

Florida State University  
Department of Statistics  
Tallahassee, Florida 32306  
Attn: Dr. Ralph A. Bradley

Princeton University  
Department of Mathematics  
Princeton, New Jersey  
Attn: Prof. J. Tukey

Columbia University  
Department of Mathematical Statistics  
New York 27, New York 10027  
Attn: Prof. T. W. Anderson

University of California  
Department of Statistics  
Berkeley 4, California  
Attn: Prof. J. Neyman

University of Washington  
Department of Mathematics  
Seattle 5, Washington  
Attn: Prof. Z. W. Birnbaum

Cornell University  
Department of Mathematics  
Ithaca, New York  
Attn: Prof. J. Wolfowitz

Harvard University  
Department of Statistics  
Cambridge, Massachusetts  
Attn: Prof. W. G. Cochran

Institute of Mathematical Statistics  
University of Copenhagen  
Copenhagen, Denmark  
Attn: Prof. Anders Hald

Columbia University  
Department of Industrial Engineering  
New York 27, New York 10027  
Attn: Prof. Cyrus Derman

Columbia University  
Department of Mathematics  
New York 27, New York 10027  
Attn: Prof. H. Robbins

New York University  
Institute of Mathematical Sciences  
New York 3, New York  
Attn: Prof. W. M. Hirsch

University of North Carolina  
Statistics Department  
Chapel Hill, North Carolina  
Attn: Prof. Walter L. Smith

Michigan State University  
Department of Statistics  
East Lansing, Michigan  
Attn: Prof. Herman Rubin

University of California, San Diego  
Department of Mathematics  
P. O. Box 109  
La Jolla, California 92038  
Attn: Prof. M. Rosenblatt

New York University  
Department of Industrial Engineering  
and Operations Research  
Bronx 63, New York  
Attn: Prof. J. H. Kao

University of Wisconsin  
Department of Statistics  
Madison, Wisconsin  
Attn: Prof. G. E. P. Box

The Research Triangle Institute  
Statistics Research Division  
505 West Chapel Hill Street  
Durham, North Carolina  
Attn: Dr. Malcolm R. Leadbetter

Florida State University  
Department of Statistics  
Tallahassee, Florida 32306  
Attn: Prof. I. R. Savage

Massachusetts Institute of Technology  
Department of Electrical Engineering  
Cambridge, Massachusetts 02139  
Attn: Dr. R. A. Howard

The Johns Hopkins University  
Department of Mathematical Statistics  
34th & Charles Streets  
Baltimore 18, Maryland  
Attn: Prof. Geoffrey S. Watson

Stanford University  
Department of Statistics  
Stanford, California 94305  
Attn: Prof. E. Parzen

Arcon Corporation  
803 Massachusetts Avenue  
Lexington 73, Massachusetts 02173  
Attn: Dr. Arthur Albert

University of California  
College of Engineering  
Operations Research Center  
Berkeley, California  
Attn: Prof. R. E. Barlow

Michigan State University  
Department of Statistics  
East Lansing, Michigan  
Attn: Prof. J. Gani

Yale University  
Department of Statistics  
New Haven, Connecticut 06520  
Attn: Prof L. J. Savage

Rocketdyne - A Division of North  
American Aviation, Inc.  
6633 Canoga Avenue  
Canoga Park, California  
Attn: Dr. N. R. Goodman  
Dr. J. M. Zimmerman

Rutgers - The State University  
Statistics Center  
New Brunswick, New Jersey 08903  
Attn: Prof. H. E. Dodge

Yale University  
Department of Statistics  
New Haven, Connecticut 06520  
Attn: Prof. F. J. Anscombe

Purdue University  
Division of Mathematical Sciences  
Lafayette, Indiana 47907  
Attn: Prof. S. S. Gupta

Cornell University  
Department of Industrial Engineering  
Ithaca, New York  
Attn: Prof. Robert Bechhofer

Stanford University  
Department of Statistics  
Stanford, California 94305  
Attn: Prof. C. Stein

Stanford University  
Department of Statistics  
Stanford, California 94305  
Attn: Prof. H. Chernoff

Mr. John E. Condon  
 Director, Office of Reliability  
 and Quality Assurance, Code KR  
 NASA Headquarters  
 400 Maryland Avenue, S. W.  
 Washington, D. C. 20546 (2 copies)

Mr. Dwight C. Cain (MGR)  
 Office of Manned Space Flight  
 Gemini Program Office  
 NASA Headquarters  
 400 Maryland Avenue, S. W.  
 Washington, D. C. 20546

Dr. John M. Walker (RET)  
 Office of Advanced Research  
 and Technology  
 NASA Headquarters  
 400 Maryland Avenue, S. W.  
 Washington, D. C. 20546

Mr. James O. Spriggs (SA)  
 Office of Space Science  
 and Applications  
 NASA Headquarters  
 400 Maryland Avenue, S. W.  
 Washington, D. C. 20546

Mr. John V. Foster, Chief  
 Systems Engineering Division  
 NASA Ames Research Center  
 Moffett Field, California 94035

Mr. Richard B. Cox, Head  
 Quality Assurance Office  
 NASA Flight Research Center  
 P. O. Box 273  
 Edwards, California 93523

Mr. Clinton T. Johnson  
 Reliability Analysis  
 NASA Flight Research Center  
 P. O. Box 273  
 Edwards, California 93523

Mr. G. Kambouris (232)  
 Office of Technical Services  
 NASA Goddard Space Flight Center  
 Greenbelt, Maryland 20771

Dr. William Wolman (600)  
 Office of Space Science  
 and Satellite Applications  
 NASA Goddard Space Flight Center  
 Greenbelt, Maryland 20771

Mr. Brooks T. Morris, Chief  
 Quality Assurance and Reliability  
 Office  
 Jet Propulsion Laboratory  
 4800 Oak Grove Drive  
 Pasadena, California 91103

Mr. Eugene H. Britt  
 Reliability & Quality  
 Assurance Officer  
 NASA Langley Research Center  
 Langley Station  
 Hampton, Virginia 23365

Mr. T. J. Edwards (SR)  
 NASA Manned Spacecraft Center  
 2101 Webster-Seabrook Road  
 Houston, Texas 77058

Mr. Robert L. Chandler, (R-QUAL-R)  
 NASA Marshall Space Flight Center  
 Huntsville, Alabama 35812

Mr. James C. Taylor (R-ASTR-R)  
 Astrionics Laboratory  
 NASA Marshall Space Flight Center  
 Huntsville, Alabama 35812

Mr. J. E. Stitt (12.000)  
 Instrument Research Division  
 NASA Langley Research Center  
 Langley Station  
 Hampton, Virginia 23365

Mr. Roy E. Currie, Jr., (R-ASTR-TN)  
 Astrionics Laboratory  
 NASA Marshall Space Flight Center  
 Huntsville, Alabama 35812

Mr. Fred L. Niemann  
 Assistant Director for Technical Programs  
 NASA North Eastern Office  
 30 Memorial Drive  
 Cambridge, Massachusetts 02142

Mr. J. Brad Aaron  
NASA Wallops Station  
Wallops Island, Virginia 23337

Mr. Edward M. James, Jr., Chief  
Technical Division  
NASA Western Operations Office  
150 Pico Boulevard  
Santa Monica, California 90406

Mr. Samuel Perrone, Assistant Chief  
Office of Reliability and  
Quality Assurance  
NASA Lewis Research Center  
21000 Brookpark Road  
Cleveland, Ohio 44135

Mr. R. Body (PA-6)  
Chief, Apollo Reliability & Quality  
Assurance Office  
NASA J. F. Kennedy Space Center  
Cocoa Beach, Florida

Mr. George A. Lemke, Director  
Apollo Reliability and Quality  
Assurance Office (MAR)  
Office of Manned Space Flight  
NASA Headquarters  
Washington, D. C. 20546 (2 copies)

Research Triangle Institute  
P. O. Box 490  
Durham, North Carolina 27702  
Attn: Project Officer NASr-40

Dr. Albert J. Kelley (RE-TG)  
Office of Advanced Research  
and Technology  
NASA Headquarters  
400 Maryland Avenue, S. W.  
Washington, D. C. 20546

Mr. I. B. Altman  
Office, Secretary of Defense (I&L)  
Quality Control and Reliability  
Room 1E815, Pentagon  
Washington, D. C. 20301

Mr. Joseph Carroll  
Bureau of Naval Weapons  
Quality Evaluation Operations  
Code FQA0-3  
Naval Weapons Plant  
Washington 25, D. C.

Mr. O. A. Cocca  
HQ, Air Force Logistics Command (MCPKQ)  
Wright-Patterson Air Force Base  
Ohio

Mr. James Coffin  
Air Force System Command (SCKAQ)  
Andrews Air Force Base  
Washington 25, D. C.

Mr. John Condon  
Director, Office of Reliability  
and Quality Assurance  
NASA Headquarters, Code KR  
Washington, D. C. 20546

Mr. Walter Foster  
Fort Detrick  
Frederick, Maryland

Mr. Fred Frishman  
OCD, Department of the Army  
419 3045 Columbia Pike  
Arlington, Virginia

Mr. Earl K. Yost  
Defense Supply Agency  
Cameron Station  
Alexandria, Virginia

Dr. Frank Grubbs  
Aberdeen Proving Grounds  
Aberdeen, Maryland

Mr. Silas Williams, Jr.  
HQ, U.S. Army Supply and  
Maintenance Command  
Chief, Quality Assurance, AMSSM-QA  
Washington, D. C. 20315

Mr. Richard Hussey  
Air Force Director of  
Procurement Management  
Room 4C253, Pentagon  
Washington, D. C. 20301

Mr. B. Kurkjian  
Harry Diamond Laboratory  
Department of the Army  
Washington 25, D. C.

Mr. Seymour Lorber  
HQ, Army Materiel Command  
Department of the Army  
Washington 25, D. C.

Mr. Joseph Mandelson  
Directorate of Quality Assurance  
Edgewood Arsenal  
Edgewood, Maryland

Mr. Arthur Marthens  
Bureau of Ships  
Department of the Navy  
Room 3012, Main Navy Bldg.  
Washington, D. C. 20360

Major John F. Mosher  
Air Force System Command (SCSVE)  
Andrews Air Force Base  
Washington 25, D. C.

Dr. W. R. Pabst, Jr.  
Chief Statistician  
Bureau of Naval Weapons (P-130)  
Washington, D. C. 20360

Mr. T. M. Vining  
HQ, Defense Supply Agency  
Cameron Station  
Alexandria, Virginia

Mr. J. Weinstein  
Chief, Mathematics Division  
Institute for Experimental Research  
U.S. Army Electronics R&D Lab.  
Fort Monmouth, New Jersey

U.S. Naval Ordnance Test Station  
China Lake, California 93557  
Attn: Statistics Branch (Code 5077)

Commanding Officer  
U.S. Naval Underwater Ordnance Station  
Newport, Rhode Island 02844  
Attn: Library (Cm4b)

Commanding Officer  
U.S. Naval Torpedo Station  
Keyport, Washington  
Code 333  
Code 334 (QEL Library)

Library  
Technical Reports Section  
U.S. Naval Postgraduate School  
Monterey, California 93940

Commander  
Philadelphia Naval Shipyard  
(Naval Boiler and Turbine Laboratory)  
Naval Base, Philadelphia, Pennsylvania 19112

Officer-in-Charge  
U.S. Naval Underwater Weapons Systems  
Engineering Center  
Newport, Rhode Island 02844  
Attn: Quality Evaluation Laboratory (QES)

Commanding Officer and Director  
U.S. Navy Electronics Laboratory (Library)  
San Diego, California 92152

Commanding Officer and Director  
U.S. Navy Mine Defense Laboratory  
Technical Library  
Panama City, Florida 32402

Commanding Officer and Director (Code 904)  
U.S. Navy Underwater Sound Laboratory  
Fort Trumbull, New London, Conn. 06321

Commanding Officer  
U.S. Naval Avionics Facility  
Indianapolis, Indiana 46218  
Attn: Library

Commanding Officer  
U.S. Naval Ammunition Depot (QETM)  
Crane, Indiana

Commanding Officer  
U.S. Naval Ammunition Depot (QEL)  
St. Juliens Creek  
Portsmouth, Virginia 23702

Commanding Officer  
U.S. Naval Weapons Station  
Seal Beach, California 90740  
Attn: Technical Library, Bldg AO-2

Officer-in-Charge  
U.S. Naval Fleet Missile Systems  
Analysis and Evaluation Group  
Corona, California

U.S. Naval Fleet Missile Systems  
Analysis and Evaluation Group  
Corona, California 91720

Commander  
U.S. Naval Ordnance Laboratory  
White Oak  
Silver Spring, Maryland  
Code (NS Div)  
Code (R - Dept)

Commanding Officer  
U.S. Naval Weapons Station (QEW)  
Concord, California

Officer-in-Charge  
U.S. Naval Mine Engineering Facility  
U.S. Naval Weapons Station  
Yorktown, Virginia - 23491

Commanding Officer  
U.S. Naval Propellant Plant  
Indian Head, Maryland  
Code (QA)  
Code (REM)

Commander  
U.S. Naval Weapons Laboratory  
Dahlgren, Virginia  
Attn: Technical Library

Chief, Bureau of Naval Weapons  
Department of the Navy  
Washington, D. C. 20360  
Code (FQ)  
Code (PQC)  
Code (P-13)  
Code (RAAV-33)

Chief of Naval Material (MAT-254)  
Department of the Navy  
Room 3025, Munitions Bldg.  
Washington, D. C. 20360

Director  
U.S. Naval Weapons Quality Assurance Office  
Quality Analysis Department  
Washington Navy Yard  
Washington, D. C. 20390

Chief of Naval Operations (Op-03EG)  
Operations Evaluation Group  
The Pentagon  
Washington, D. C. 20350

Commanding Officer (ADLR)  
U. S. Naval Air Development Center  
Johnsville, Pa.  
Attn: NADC Library

Commander  
U.S. Naval Missile Center  
Point Mugu, California  
Attn: Technical Library, Code NO322

3 copies

Commanding Officer  
U.S. Naval Construction Battalion Center  
Gulfport, Mississippi

U.S. Naval Construction Battalion Center  
Port Hueneme, California 93041  
Code 92

U.S. Naval Ammunition Depot  
Navy No. 66, c/o FPO  
San Francisco, California 96612  
Attn: Weapons Technical Library

2 copies

Commander, U.S. N.O.T.S.  
Pasadena Annex  
3202 East Foothill Blvd.  
Pasadena, California 91107  
Attn: Pasadena Annex Library



UNCLASSIFIED

Security Classification

## DOCUMENT CONTROL DATA - R&amp;D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author)		2a. REPORT SECURITY CLASSIFICATION	
University of California, Berkeley		Unclassified	
		2b. GROUP	
3. REPORT TITLE			
Exponential Life Test Procedures When the Distribution has Monotone Failure Rate			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)			
Technical Report, October 1964			
5. AUTHOR(S) (Last name, first name, initial)			
Barlow, Richard E. and Proschan, Frank			
6. REPORT DATE		7a. TOTAL NO. OF PAGES	7b. NO. OF REFS
November 1964		34	22
8a. CONTRACT OR GRANT NO.		9a. ORIGINATOR'S REPORT NUMBER(S)	
Nomr-3656(18)		ORC 64-29	
b. PROJECT NO.			
c.		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
d.			
10. AVAILABILITY/LIMITATION NOTES			
Releasable without limitations on dissemination.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY	
		Logistics and Mathematical Statistics Branch, Office of Naval Research Washington, D.C. 20360	
13. ABSTRACT			
<p>This paper studies the performance of the standard life test procedures developed for the exponential distribution when the underlying distribution actually has an increasing (decreasing) failure rate. The standard exponential life test procedures are shown to yield estimates of the unknown mean biased on the high side in the increasing failure rate case. The practical implications of these results are that the consumer tends to accept unduly often poor quality lots, when the producer has used an exponential life test procedure and the underlying failure distribution actually had increasing failure rate.</p> <p>In addition a number of theoretical results are obtained concerning order statistics and their spacings from distributions with increasing failure rate. These results are useful in the study of the life test procedures and also are of independent interest.</p>			

14	KEY WORDS	LINK A		LINK B		LINK C	
		ROLE	WT	ROLE	WT	ROLE	WT
<p>Life Testing</p> <p>Monotone Failure Rate</p> <p>Exponential estimates</p> <p>Bias of the mean life</p>							

INSTRUCTIONS

1. **ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (*corporate author*) issuing the report.

2a. **REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

2b. **GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

3. **REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parenthesis immediately following the title.

4. **DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

5. **AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

6. **REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

7a. **TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

7b. **NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

8a. **CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

8b, 8c, & 8d. **PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

9a. **ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

9b. **OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (*either by the originator or by the sponsor*), also enter this number(s).

10. **AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through \_\_\_\_\_."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through \_\_\_\_\_."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through \_\_\_\_\_."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

11. **SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

12. **SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address.

13. **ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

14. **KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical content. The assignment of links, roles, and weights is optional.